

Goat

Purina® Goat Nutrition Program

Purina[®] Goat Grower 16 pq.0015

Complete feed for growth, development and maintenance of goats

Purina® Goat Grower 16 DQ.0015 is a pelleted complete feed formulated for the optimum growth, development and maintenance of goats. Whether you raise meat goats or dairy goats, Purina® Goat Grower 16 DQ.0015 feed is designed to meet their exacting needs.

Purina® products deliver the nutrition and performance you expect. Try Purina® Goat Grower 16 DQ.0015 feed and see the difference it can make in your goats' overall growth and health.



FEATURE	BENEFIT
Nutritionally Complete	Helps provides the proper balance of high-quality proteins, vitamins, minerals and other nutrients Built in roughage source helps minimize clean-up from wasted or uneaten forage
Palatable, Premium Ingredients	Consistent quality helps assure high palatability ans supports performance
Pelleted	Helps reduce sorting, easy to handle
Urinary Acidifiers including Ammonium Chloride and/or Ammonium Sulfate	Helps reduce the incidence of urinary calculi
Options	Non-medicated option available (depending on plant)





Goat

Purina® Goat Nutrition Program

Purina[®] Goat Grower 16 pq.0015



Guaranteed Analysis:

ACTIVE DRUG INGREDIENTS

Decoquinate	0.0015 %
	(6.8 mg/lb)

NUTRIENT	LEVEL	
Crude Protein, (Min)	16.00%	
This includes not more than 1.5% equivalent crude protein from non-protein nitrogen		
Crude Fat, (min)	2.00%	
Crude Fiber, (max)	16.00%	
Acid Detergent Fiber (ADF) (max)	22.00%	
Calcium (Ca), (min)	0.75%	
Calcium (Ca), (max)	1.25%	
Phosphorus (P), (min)	0.50%	
Salt (NaCl), (min)	0.50%	
Salt (NaCl), (max)	1.00%	
Copper (Cu), (min)	22.00 ppm	
Copper (Cu), (max)	27.00 ppm	
Selenium (Se), (min)	0.30 ppm	
Vitamin A, (min)	5,000 IU/lb	

Ingredients:

Processed Grain By-Products, Grain Products, Roughage Products, Plant Protein Products, Molasses Products, Forage Products, Calcium Carbonate, Salt, Ammonium Chloride, Lignin Sulfonate, Colored with Iron Oxide, Dicalcium Phosphate, Monocalcium Phosphate, Sodium Selenite, Vitamin D₃ Supplement, Vitamin E Supplement, Vitamin A Supplement, Zinc Oxide, Cobalt Carbonate, Manganese Sulfate, Ethylenediamine Dihydroiodide, Zinc Sulfate, Basic Copper Chloride.

Directions:

Feed at a rate of 1.67 lb to goats/kids weighing 50 lb to provide 22.7mg decoquinate per 100 lb of bodyweight (0.5mg/kg) per day. Feed at least 28 days during time of exposure to coccidiosis or when it is likely to be a hazard.

Warning: Do not feed to goats producing milk for food.

Important:

- Start kids on colostrum milk from birth to 3 days of age; feed Land O'Lakes® Doe's Match® Kid Milk Replacer from 2 days until 4-8 weeks of age.
- If kids have not been creep fed or they were purchased, then they should be placed on grass hay on a free-choice basis after arrival.
- 3. When making a ration change, allow 7 to 10 days for animals to adjust to the new ration.
- 4. Keep a constant supply of complete feed available. Do not let fine material accumulate in feeders
- Provide adequate bunk space for each animal. Bunks should be well-protected and well-managed to prevent feed from becoming wet or moldy.
- 6. Provide a source of fresh, clean water at all times.
- 7. Feed Purina® Goat Mineral free-choice.
- 8. Consult your veterinarian for the recommended health program for your local area. This includes internal and external parasite control. In groups of goats there are certain animals that experience chronic bloat, or other digestive disturbances and consequently, are poor performers. In addition, excess feed consumption, severe weather changes resulting in erratic feed consumption and poorly managed feeding practices, can increase the incidence of bloat in all goats. If bloating occurs, the above management practices should be reviewed and your veterinarian consulted.

Caution

Store in a dry, well-ventilated area protected from rodents and insects. Do not feed moldy or insect-infested feed to animals as it may cause illness, performance loss or death.

This product contains copper and should not be fed to sheep. Bentonite should not be used in decoquinate feeds.

USE ONLY AS DIRECTED.

